

PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Summary Sort Save Text Clip Add Order

Text Version

Show: 20 Items 1-20 of 172 Page 1 of 9 Select page: 1 2 3 4 5 6 7 8 9

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journal Browser
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

- ☐ 1: [Baardsnes J, Kondejewski LH, Hodges RS, Chao H, Kay C, Davies PL.](#) Related Articles
New ice-binding face for type I antifreeze protein.
FEBS Lett. 1999 Dec 10;463(1-2):87-91.
PMID: 10601644 [PubMed - indexed for MEDLINE]
- ☐ 2: [Loewen MC, Chao H, Houston ME Jr, Baardsnes J, Hodges RS, Kay CM, Sykes BD, Sonnichsen FD, Davies PL.](#) Related Articles
Alternative roles for putative ice-binding residues in type I antifreeze protein.
Biochemistry. 1999 Apr 13;38(15):4743-9.
PMID: 10200162 [PubMed - indexed for MEDLINE]
- ☐ 3: [Cheng A, Merz KM Jr.](#) Related Articles
Ice-binding mechanism of winter flounder antifreeze proteins.
Biophys J. 1997 Dec;73(6):2851-73.
PMID: 9414201 [PubMed - indexed for MEDLINE]
- ☐ 4: [Baardsnes J, Jelokhani-Niaraki M, Kondejewski LH, Kuiper MJ, Kay CM, Hodges RS, Davies PL.](#) Related Articles
Antifreeze protein from shorthorn sculpin: identification of the ice-binding surface.
Protein Sci. 2001 Dec;10(12):2566-76.
PMID: 11714925 [PubMed - indexed for MEDLINE]
- ☐ 5: [Chao H, Houston ME Jr, Hodges RS, Kay CM, Sykes BD, Loewen MC, Davies PL, Sonnichsen FD.](#) Related Articles
A diminished role for hydrogen bonds in antifreeze protein binding to ice.
Biochemistry. 1997 Dec 2;36(48):14652-60.
PMID: 9398184 [PubMed - indexed for MEDLINE]
- ☐ 6: [Zhang W, Laursen RA.](#) Related Articles
Structure-function relationships in a type I antifreeze polypeptide. The role of threonine methyl and hydroxyl groups in antifreeze activity.
J Biol Chem. 1998 Dec 25;273(52):34806-12.
PMID: 9857006 [PubMed - indexed for MEDLINE]
- ☐ 7: [Wen D, Laursen RA.](#) Related Articles
Structure-function relationships in an antifreeze polypeptide. The effect of added bulky groups on activity.
J Biol Chem. 1993 Aug 5;268(22):16401-5.
PMID: 8344925 [PubMed - indexed for MEDLINE]
- ☐ 8: [Dalal P, Sonnichsen FD.](#) Related Articles
Source of the ice-binding specificity of antifreeze protein type I.
J Chem Inf Comput Sci. 2000 Sep-Oct;40(5):1276-84.
PMID: 11045824 [PubMed - indexed for MEDLINE]
- ☐ 9: [Houston ME Jr, Chao H, Hodges RS, Sykes BD, Kay CM, Sonnichsen FD, Loewen MC, Davies PL.](#) Related Articles

Binding of an oligopeptide to a specific plane of ice.

J Biol Chem. 1998 May 8;273(19):11714-8.

PMID: 9565593 [PubMed - indexed for MEDLINE]

- ☐ **10:** Lin Q, Ewart KV, Yan Q, Wong WK, Yang DS, Hew CL. Related Articles
Secretory expression and site-directed mutagenesis studies of the winter flounder skin-type antifreeze polypeptides.
Eur J Biochem. 1999 Aug;264(1):49-54.
PMID: 10447672 [PubMed - indexed for MEDLINE]
- ☐ **11:** Graether SP, DeLuca CI, Baardsnes J, Hill GA, Davies PL, Jia Z. Related Articles, Protein, Structure
Quantitative and qualitative analysis of type III antifreeze protein structure and function.
J Biol Chem. 1999 Apr 23;274(17):11842-7.
PMID: 10207002 [PubMed - indexed for MEDLINE]
- ☐ **12:** Sicheri F, Yang DS. Related Articles, Protein
Ice-binding structure and mechanism of an antifreeze protein from winter flounder.
Nature. 1995 Jun 1;375(6530):427-31.
PMID: 7760940 [PubMed - indexed for MEDLINE]
- ☐ **13:** Jia Z, DeLuca CI, Chao H, Davies PL. Related Articles, Protein, Structure
Structural basis for the binding of a globular antifreeze protein to ice.
Nature. 1996 Nov 21;384(6606):285-8.
PMID: 8918883 [PubMed - indexed for MEDLINE]
- ☐ **14:** Yang DS, Hon WC, Bubanko S, Xue Y, Seetharaman J, Hew CL, Sicheri F. Related Articles
Identification of the ice-binding surface on a type III antifreeze protein with a "flatness function" algorithm.
Biophys J. 1998 May;74(5):2142-51.
PMID: 9591641 [PubMed - indexed for MEDLINE]
- ☐ **15:** Harding MM, Ward LG, Haymet AD. Related Articles
Type I 'antifreeze' proteins. Structure-activity studies and mechanisms of ice growth inhibition.
Eur J Biochem. 1999 Sep;264(3):653-65. Review.
PMID: 10491111 [PubMed - indexed for MEDLINE]
- ☐ **16:** Chao H, Hodges RS, Kay CM, Gauthier SY, Davies PL. Related Articles
A natural variant of type I antifreeze protein with four ice-binding repeats is a particularly potent antifreeze.
Protein Sci. 1996 Jun;5(6):1150-6.
PMID: 8762146 [PubMed - indexed for MEDLINE]
- ☐ **17:** Wen D, Laursen RA. Related Articles
A model for binding of an antifreeze polypeptide to ice.
Biophys J. 1992 Dec;63(6):1659-62.
PMID: 1489916 [PubMed - indexed for MEDLINE]
- ☐ **18:** Graether SP, Kuiper MJ, Gagne SM, Walker VK, Jia Z, Sykes BD, Davies PL. Related Articles, Protein, Structure
Beta-helix structure and ice-binding properties of a hyperactive antifreeze protein from an insect.
Nature. 2000 Jul 20;406(6793):325-8.
PMID: 10917537 [PubMed - indexed for MEDLINE]
- ☐ **19:** Liou YC, Tocilj A, Davies PL, Jia Z. Related Articles, Protein, Structure
Mimicry of ice structure by surface hydroxyls and water of a beta-helix antifreeze protein.

☐ 20: [DeLuca CI, Chao H, Sonnichsen FD, Sykes BD, Davies PL.](#)

[Related Articles](#)

Effect of type III antifreeze protein dilution and mutation on the growth inhibition of ice.


Biophys J. 1996 Nov;71(5):2346-55.

PMID: 8913575 [PubMed - indexed for MEDLINE]

Show: Items 1-20 of 172 Page 1 of 9 Select page: [1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#)

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

i686-pc-linux-gnu Jul 16 2002 16:34:53

[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[PopSet](#)[Taxonomy](#)[OMIM](#)[Books](#)

Search

PubMed



for anti-freeze AND protein

Go

Clear

[Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)[About Entrez](#)

Display

Summary



Sort



Save

Text

Clip Add

Order

Text Version

Show:

20

Items 1-11 of 11

One page.

Entrez PubMed

[Overview](#)[Help | FAQ](#)[Tutorial](#)[New/Noteworthy](#)[E-Utilities](#)

PubMed Services

[Journal Browser](#)[MeSH Browser](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[LinkOut](#)[Cubby](#)

Related Resources

[Order Documents](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)[Privacy Policy](#)

- ☐
- 1:
- [Baderschneider B, Crevel RW, Earl LK, Lalljie A, Sanders DJ, Sanders IJ.](#)

[Related Articles](#)

Sequence analysis and resistance to pepsin hydrolysis as part of an assessment of the potential allergenicity of ice structuring protein type III HPLC 12.

Food Chem Toxicol. 2002 Jul;40(7):965-78.

PMID: 12065219 [PubMed - in process]

- ☐
- 2:
- [Barrett J.](#)

[Related Articles](#)

Thermal hysteresis proteins.

Int J Biochem Cell Biol. 2001 Feb;33(2):105-17. Review.

PMID: 11240367 [PubMed - indexed for MEDLINE]

- ☐
- 3:
- [Lin Q, Ewart KV, Yang DS, Hew CL.](#)

[Related Articles](#)

Studies of a putative ice-binding motif in winter flounder skin-type anti-freeze polypeptide.

FEBS Lett. 1999 Jun 25;453(3):331-4.

PMID: 10405170 [PubMed - indexed for MEDLINE]

- ☐
- 4:
- [Hey JM, MacFarlane DR.](#)

[Related Articles](#)

Crystallization of ice in aqueous solutions of glycerol and dimethyl sulfoxide 2: ice crystal growth kinetics.

Cryobiology. 1998 Sep;37(2):119-30.

PMID: 9769162 [PubMed - indexed for MEDLINE]

- ☐
- 5:
- [Lagneaux D, Huhtinen M, Koskinen E, Palmer E.](#)

[Related Articles](#)

Effect of anti-freeze protein (AFP) on the cooling and freezing of equine embryos as measured by DAPI-staining.

Equine Vet J Suppl. 1997 Dec;(25):85-7.

PMID: 9593536 [PubMed - indexed for MEDLINE]

- ☐
- 6:
- [Chandrasekaran EV, Jain RK, Vig R, Matta KL.](#)

[Related Articles](#)

The enzymatic sulfation of glycoprotein carbohydrate units: blood group T-hapten specific and two other distinct Gal:3-O-sulfotransferases as evident from specificities and kinetics and the influence of sulfate and fucose residues occurring in the carbohydrate chain on C-3 sulfation of terminal Gal.

Glycobiology. 1997 Sep;7(6):753-68.

PMID: 9376678 [PubMed - indexed for MEDLINE]

- ☐
- 7:
- [Dautel H, Knulle W.](#)

[Related Articles](#)

The supercooling ability of ticks (Acari, Ixodoidea).

J Comp Physiol [B]. 1996;166(8):517-24.

PMID: 8981761 [PubMed - indexed for MEDLINE]

- ☐
- 8:
- [Wen D, Laursen RA.](#)

[Related Articles](#)

Structure-function relationships in an antifreeze polypeptide. The role of neutral, polar amino acids.

he

☐ **9:** [Lange R, Larroque C, Anzenbacher P.](#)

[Related Articles](#)

The cholesterol-side-chain-cleaving cytochrome P450 spin-state equilibrium. 1.
Thermodynamic analysis.
Eur J Biochem. 1992 Jul 1;207(1):69-73.
PMID: 1628662 [PubMed - indexed for MEDLINE]

☐ **10:** [Kelm S, Schauer R.](#)

[Related Articles](#)

The galactose-recognizing system of rat peritoneal macrophages; identification and characterization of the receptor molecule.
Biol Chem Hoppe Seyler. 1988 Aug;369(8):693-704.
PMID: 2850817 [PubMed - indexed for MEDLINE]

☐ **11:** [Knight CA, DeVries AL, Oolman LD.](#)

[Related Articles](#)

Fish antifreeze protein and the freezing and recrystallization of ice.
Nature. 1984 Mar 15-21;308(5956):295-6.
PMID: 6700733 [PubMed - indexed for MEDLINE]

Display	Summary	Sort	Save	Text	Clip Add	Order
---------	---------	------	------	------	----------	-------

Show: 20 Items 1-11 of 11 One page.

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Search for

[Limits](#)
[Preview/Index](#)
[History](#)
[Clipboard](#)
[Details](#)

[About Entrez](#)

Text Version

[Entrez PubMed](#)
[Overview](#)
[Help | FAQ](#)
[Tutorial](#)
[New/Noteworthy](#)
[E-Utilities](#)

[PubMed Services](#)
[Journal Browser](#)
[MeSH Browser](#)
[Single Citation Matcher](#)
[Batch Citation Matcher](#)
[Clinical Queries](#)
[LinkOut](#)
[Cubby](#)

[Related Resources](#)
[Order Documents](#)
[NLM Gateway](#)
[TOXNET](#)
[Consumer Health](#)
[Clinical Alerts](#)
[ClinicalTrials.gov](#)
[PubMed Central](#)

[Privacy Policy](#)

☐ 1: Int J Biochem Cell Biol 2001 Feb;33(2):105-17

[Related Articles, Books, LinkOut](#)

ELSEVIER SCIENCE
 FULL-TEXT ARTICLE

Thermal hysteresis proteins.

Barrett J.

Institute of Biological Sciences, University of Wales, Aberystwyth, Penglais,
 Ceredigion SY23 3DA, Aberystwyth, UK. jzb@aber.ac.uk


Extreme environments present a wealth of biochemical adaptations. Thermal hysteresis proteins (THPs) have been found in vertebrates, invertebrates, plants, bacteria and fungi and are able to depress the freezing point of water (in the presence of ice crystals) in a non-colligative manner by binding to the surface of nascent ice crystals. The THPs comprise a disparate group of proteins with a variety of tertiary structures and often no common sequence similarities or structural motifs. Different THPs bind to different faces of the ice crystal, and no single mechanism has been proposed to account for THP ice binding affinity and specificity. Experimentally THPs have been used in the cryopreservation of tissues and cells and to induce cold tolerance in freeze susceptible organisms. THPs represent a remarkable example of parallel and convergent evolution with different proteins being adapted for an anti-freeze role.

Publication Types:

- Review
- Review, Academic

PMID: 11240367 [PubMed - indexed for MEDLINE]

[Write to the Help Desk](#)
[NCBI | NLM | NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act | Disclaimer](#)

[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[PopSet](#)[Taxonomy](#)[OMIM](#)[Books](#)Search

for

Go

Clear

[Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)[About Entrez](#)

Display

Abstract

Sort

Save

Text

Clip Add

Order

[Text Version](#)[Entrez PubMed](#)[Overview](#)[Help | FAQ](#)[Tutorial](#)[New/Noteworthy](#)[E-Utilities](#)[PubMed Services](#)[Journal Browser](#)[MeSH Browser](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[LinkOut](#)[Cubby](#)[Related Resources](#)[Order Documents](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)[Privacy Policy](#)☐ 1: Annu Rev Physiol 2001;63:327-57[Related Articles, Books, LinkOut](#)Full text article at
physiol.AnnualReviews.org

Antifreeze and ice nucleator proteins in terrestrial arthropods.

Duman JG.Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana 46556, USA. duman.1@nd.edu

Terrestrial arthropods survive subzero temperatures by becoming either freeze tolerant (survive body fluid freezing) or freeze avoiding (prevent body fluid freezing). Protein ice nucleators (PINs), which limit supercooling and induce freezing, and antifreeze proteins (AFPs), which function to prevent freezing, can have roles in both freeze tolerance and avoidance. Many freeze-tolerant insects produce hemolymph PINs, which induce freezing at high subzero temperatures thereby inhibiting lethal intracellular freezing. Some freeze-tolerant species have AFPs that function as cryoprotectants to prevent freeze damage. Although the mechanism of this cryoprotection is not known, it may involve recrystallization inhibition and perhaps stabilization of the cell membrane. Freeze-avoiding species must prevent inoculative freezing initiated by external ice across the cuticle and extend supercooling abilities. Some insects remove PINs in the winter to promote supercooling, whereas others have selected against surfaces with ice-nucleating abilities on an evolutionary time scale. However, many freeze-avoiding species do have proteins with ice-nucleating activity, and these proteins must be masked in winter. In the beetle *Dendroides canadensis*, AFPs in the hemolymph and gut inhibit ice nucleators. Also, hemolymph AFPs and those associated with the layer of epidermal cells under the cuticle inhibit inoculative freezing. Two different insect AFPs have been characterized. One type from the beetles *D. canadensis* and *Tenebrio molitor* consists of 12- and 13-mer repeating units with disulfide bridges occurring at least every six residues. The spruce budworm AFP lacks regular repeat units. Both have much higher activities than any known AFPs.

Publication Types:

- Review
- Review Literature

PMID: 11181959 [PubMed - indexed for MEDLINE]

Display

Abstract

Sort


Save

Text

Clip Add

Order

[Write to the Help Desk](#)

[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[PopSet](#)[Taxonomy](#)[OMIM](#)[Books](#)Search **PubMed** for[Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)[About Entrez](#)[Display](#)[Abstract](#)[Sort](#)[Save](#)[Text](#)[Clip Add](#)[Order](#)[Text Version](#)[Entrez PubMed](#)[Overview](#)[Help | FAQ](#)[Tutorial](#)[New/Noteworthy](#)[E-Utilities](#)[PubMed Services](#)[Journal Browser](#)[MeSH Browser](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[LinkOut](#)[Cubby](#)[Related Resources](#)[Order Documents](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)[Privacy Policy](#)☐ 1: Protein Sci 1996 Jun;5(6):1150-6[Related Articles, Books, LinkOut](#)

A natural variant of type I antifreeze protein with four ice-binding repeats is a particularly potent antifreeze.

Chao H, Hodges RS, Kay CM, Gauthier SY, Davies PL.

Protein Engineering Network of Centres of Excellence, University of Alberta, Edmonton, Canada.

A 4.3-kDa variant of Type I antifreeze protein (AFP9) was purified from winter flounder serum by size exclusion chromatography and reversed-phase HPLC. By the criteria of mass, amino acid composition, and N-terminal sequences of tryptic peptides, this variant is the posttranslationally modified product of the previously characterized AFP gene 21a. It has 52 amino acids and contains four 11-amino acid repeats, one more than the major serum AFP components. The larger protein is completely alpha-helical at 0 degree C, with a melting temperature of 18 degrees C. It is considerably more active as an antifreeze than the three-repeat winter flounder AFP and the four-repeat yellowtail flounder AFP, both on a molar and a mg/mL basis. Several structural features of the four-repeat winter flounder AFP, including its larger size, additional ice-binding residues, and differences in ice-binding motifs might contribute to its greater activity. Its abundance in flounder serum, together with its potency as an antifreeze, suggest that AFP9 makes a significant contribution to the overall freezing point depression of the host.

PMID: 8762146 [PubMed - indexed for MEDLINE]

[Display](#)[Abstract](#)[Sort](#)[Save](#)[Text](#)[Clip Add](#)[Order](#)[Write to the Help Desk](#)[NCBI | NLM | NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act | Disclaimer](#)

Search PubMed for

Limits Preview/Index History Clipboard Details

About Entrez

Abstract Text

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journal Browser
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

☐ 1: FEBS Lett 1999 Dec 10;463(1-2):87-91

Related Articles, Books, LinkOut

ELSEVIER SCIENCE
FULL-TEXT ARTICLE

New ice-binding face for type I antifreeze protein.

Baardsnes J, Kondejewski LH, Hodges RS, Chao H, Kay C, Davies PL.

Department of Biochemistry and the Protein Engineering Network of Centres of Excellence, Queen's University, Kingston, Ont., Canada.

Type I antifreeze protein (AFP) from winter flounder is an alanine-rich, 37 amino acid, single alpha-helix that contains three 11 amino acid repeats (Thr-X(2)-Asx-X(7)), where X is generally Ala. The regularly spaced Thr, Asx and Leu residues lie on one face of the helix and have traditionally been thought to form hydrogen bonds and van der Waals interactions with the ice surface. Recently, substitution experiments have called into question the importance of Leu and Asn for ice-binding. Sequence alignments of five type I AFP isoforms show that Leu and Asn are not well conserved, whereas Ala residues adjacent to the Thr, at right angles to the Leu/Asn-rich face, are completely conserved. To investigate the role of these Ala residues, a series of Ala to Leu steric mutations was made at various points around the helix. All the substituted peptides were fully alpha-helical and remained as monomers in solution. Wild-type activity was retained in A19L and A20L. A17L, where the substitution lies adjacent to the Thr-rich face, had no detectable antifreeze activity. The nearby A21L substitution had 10% wild-type activity and demonstrated weak interactions with the ice surface. We propose a new ice-binding face for type I AFP that encompasses the conserved Ala-rich surface and adjacent Thr.

PMID: 10601644 [PubMed - indexed for MEDLINE]

Abstract Text

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer